OXFORD DICTIONARY OF

BIOCHEMISTRY AND MOLECULAR BIOLOGY

removing

Fully up-dated revised edition

BEST AVAILABLE COPY

OXFORD DICTIONARY OF

BIOCHEMISTRY AND MOLECULAR BIOLOGY

REVISED EDITION

Managing Editor Dr A D Smith University College London

General Editors Professor S P Datta University College London

Dr G H Smith University College London

Professor P N Campbell (Chairman) University College London

Dr R Bentley University of Pittsburgh

Dr H A McKenzie Australian Defence Force Ac.

Subject Editors Dr D A Bender University College London

Dr A J Harris University of Queensland

Professor T W Goodwin University of Liverpool

Dr H A McKenzie Australian Defence Force Ac.

Dr J H Parish University of Leeds

Dr C Stanford University College London

OXFORD UNIVERSITY PRESS

BEST AVAILABLE COPY

OXFORD

INTUEDSITY DRESS

Great Clarendon Street, Oxford 0x2 6DP

Oxford University Press is a department of the University of Oxford. It furthers the University's objective of excellence in research, scholarship, and education by publishing worldwide in

Oxford New York

Athens Auckland Bangkok Bogotá Buenos Aires Cape Town Chennai Dar es Salaam Delhi Florence Hong Kong Istanbul Karachi Kolkata Kuala Lumpur Madrid Melbourne Mexico City Mumbai Nairobi Paris São Paulo Shanghai Singapore Taipei Tokyo Toronto Warsaw and associated companies in Berlin Ibadan

Oxford is a registered trade mark of Oxford University Press in the UK and in certain other countries

Published in the United States by Oxford University Press Inc., New York

© The General Editors, 1997

The authors have asserted their right under the Copyright, Designs and Patents Act, 1988, to be identified as authors of this work

First published 1997 Revised edition 2000 Reprinted 2001

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, without the prior permission in writing of Oxford University Press, or as expressly permitted by law, or under terms agreed with the appropriate reprographics rights organization. Enquiries concerning reproduction outside the scope of the above should be sent to the Rights Department, Oxford University Press, at the address above

You must not circulate this book in any other binding or cover and you must impose this same condition on any acquirer

A catalogue record for this book is available from the British Library

Library of Congress Cataloging in Publication Data (Data applied for) ISBN 0 19 850673 2

> Printed in Great Britain by Butler & Tanner Ltd, Frome

biochemists nation outsid ning to revisi

y of Biocher

rovides a co

chemistry a

are short bu

information

over 17 000

nemical subs

they are inv

epts in mole

hiochemical

ernative nam sted and will

iere the inter ochemical no

de the struct

unds of inter

strations of

ontains over

which descr

the reactions

se codes to f

ence databas

rided for rele

ctionary of B

logy will ser

tograph: Invo . Private colle Reproduced by . Hepworth Es

BEST AVAILABLE COPY

components of the light scattered by scattering particles when illuminated by unpolarized light. If the scattering particles are isotropic the depolarization ratio will be zero; if they are anisotropic it will not be zero.

deproteinization or deproteinisation the process of removing protein from a biological sample, either by precipitation of protein from solution or by hydrolysis of protein with prote-

olytic enzymes.

depside any of a family of natural or artificial compounds derived from two or more molecules of various trihydroxybenzoic acids by esterification of a carboxyl group on one molecule with a hydroxyl group on another molecule (which may or may not be of the same structure as the first). By analogy with peptides, depsides derived from two, three, etc., or several trihydroxybenzoic acid units may be termed didepside, tridepside, etc., or polydepside, respectively. Depsides occur widely in lichens and also as components of some tannins.

depsidone any compound derived from two phenolcarboxylic acid molecules (which may or may not be the same molecules) that are joined together by both an ester and an ether link.

Such substances are found in lichens.

depsipeptide any linear or macrocyclic compound containing (alternating) amino-acid and hydroxy-acid residues, linked by amide, N-methylamide, and ester bonds. Among the naturally occurring cyclodepsipeptides (sometimes alternatively termed peptolides) are the antibiotics valinomycin, the enniatins, and the monamycins.

depurinate to remove purine residues from a polynucleotide.

-depurinated adj.; depurination n.

depyrimidinate to remove pyrimidine residues from a polynu-

cleotide. —depyrimidinated adj.; depyrimidination n.

derepression the phenomenon in which a repressor is prevented from interacting with the operator component of an operon and an increase in the synthesis of a gene product thereby occurs. With an inducible enzyme, the inducer derepresses the operon. Derepression may also result from mutations, either in the regulator(y) gene, thus blocking repressor synthesis, or in the operator gene, thus rendering it insensitive to the repressor.

deRib (sometimes) symbol for 2-deoxy-D-ribose (dRib is the

correct symbol).

derivative 1 any compound that may, at least theoretically, be formed from another compound to which it is structurally related. 2 see derivative of a function.

derivative of a function (in mathematics) the limit of the ratio of an increment of a dependent variable to the corresponding increment of an associated independent variable as the latter increment tends to zero.

derivatize or derivatise to convert a chemical compound into a

derivative. —derivatization or derivatisation n.

dermaseptin any of a series of antimicrobial peptides derived from amphibian skin. Dermaseptins s1-5 constitute a family of cationic lysine-rich amphipathic antifungal peptides of 28-34 residues. Presumed to protect the naked frog skin from infections, they have wider application, being the first vertebrate peptides to show lethal effects against the filamentous fungi that can cause infections in immunodeficiency syndrome, or during the use of immunosuppressive agents. Compare

dermatan sulfate or chondroitin sulfate B any of a group of glycosaminoglycans with repeats consisting of $\beta 1 \rightarrow 4$ -linked Liduronyl- $(\beta 1 \rightarrow 3)$ -N-acetyl-D-galactosamine 4-sulfate units. They are important components of ground substance or intercellular cement of skin and some connective tissues (see matrix (def. 3)). They usually occur attached to proteins as proteoglycans, the mode of linkage being the same as for chondroitin sulfate. A hereditary deficiency in the ability to degrade dermatan sulfate characterizes certain types of mucopolysaccharidosis.

dermis or corium the thick layer of tissue forming part of the skin and lying beneath the epidermis. It consists of loose connective tissue in which are blood capillaries, smooth-muscle

fibres, sweat glands and sebaceous glands with their ducts, hair follicles, and sensory nerve endings. —dermal, dermic adj. dermorphin any of a group of heptapeptide amides of the general structure H-Tyr-DAla-Phe-Gly-Tyr-Xaa-Ser-NH2, isolated from the skin of frogs of the genus Phyllomedusa and some other amphibians; they show high and long-lasting opiate-like activity.

DES abbr. for diethylstilbestrol.

des+ prefix 1 (in the trivial name of a polycyclic compound, e.g. a steroid) indicating the removal of a terminal ring, with addition of a hydrogen atom at each junction arm with the adjacent ring; it should be followed by the italic capital letter designating the terminal ring in question, e.g. des-A-androstane. 2 (in the trivial name of a polypeptide) indicating the removal of an amino-acid residue, terminal or otherwise; it should be followed by the name or symbol for the amino acid in question and an arabic number indicating its position in the normal polypeptide, e.g. des-7-proline-oxytocin, or des-Pro7oxytocin. 3 indicating the removal of a specified atom or group from a molecule of an indicated substance; see desamido+, desthio+; see also descarboxy-clotting factor. 4 (formerly) a variant of de+ (still preferred in some languages, e.g. French, German).

desalinate to remove salt, especially from seawater to render it suitable for drinking or irrigation. Compare desalt.

-desalination n.

desalt to remove small, usually inorganic ions from a sample. Methods that may be used include electrodialysis, electrophoresis, gel filtration, and ion-exchange chromatography. See also deionize. Compare desalinate.

desamidation the hydrolysis of one or more carboxamide

groups in a molecule to carboxyl groups.

desamido+ prefix indicating desamidation of a molecule, espe-

cially of a peptide.

desaturase general name for any enzyme catalysing a desaturation reaction. The name specifically often refers to any of several fatty-acyl-CoA desaturases, enzymes coming within subsubclass EC 1.14.99; in animals these are associated with the endoplasmic reticulum, require dioxygen, and insert double bonds into saturated and Z(i.e. cis)-unsaturated fatty acids. They also occur in higher plants, protozoa, and fungi. These enzymes are specific for position in the fatty-acyl chain, and are termed Δ^5 -desaturase, Δ^9 -desaturase, and so on, according to the distance from the carboxyl carbon atom to the first carbon atom of the double bond. In mammals, there is a lack of any desaturase that inserts a double bond nearer to the terminal methyl group of the fatty acid than nine carbon atoms, hence the need for n-3 and n-6 precursor fatty acids (the essential fatty acids). See also linoleic family, linolenic family.

desaturation 1 any process or reaction in which an organic compound becomes unsaturated (def. 2), e.g. by removal of two hydrogen atoms, or a hydrogen atom and a hydroxyl group, from adjacent carbon atoms. 2 any process in which ligands are removed from a macromolecule so that all the binding sites for that ligand are no longer occupied. —desaturate vb.;

desaturated adj.

descarboxy-clotting factor any of the abnormal blood clotting factors containing glutamic acid residues instead of ycarboxyglutamic-acid residues. They are formed in animals deficient in vitamin K or during administration of vitamin K antagonists. Compare PIVKA.

Descartes, René (1596-1650), French philosopher and mathe-

matician. See also Cartesian.

descending chromatography any chromatographic technique in which a liquid mobile phase runs downwards through

the supporting matrix.

descriptive name name of an organic substance that is more descriptive of action or function and often more convenient than its trivial name, semisystematic name, systematic name, or other type of name permitted by the rules of chemical nomenclature.